



GoatCare • 2X

TYPE C MEDICATED FEED

GOAT DEWORMER

Contains Rumatel® (Morantel Tartrate)

TREATS
12
FIFTY LB
GOATS



For removal and control of mature gastrointestinal nematode infections of goats including *Haemonchus contortus*, *Ostertagia (Teladorsagia) circumcincta*, and *Trichostrongylus axei*.

ACTIVE DRUG INGREDIENT
Morantel Tartrate 0.194% (0.88 grams per pound)

Rumatel is a registered trademark of Phibro Animal Health Corporation.
See Back Panel for Additional Information

NET WT: 3 LB (1.36 KG)



GoatCare•2X

TYPE C MEDICATED FEED

GOAT DEWORMER



For removal and control of mature gastrointestinal nematode infections of goats including *Haemonchus contortus*, *Ostertagia (Teladorsagia) circumcincta* and *Trichostrongylus axei*.

ACTIVE INGREDIENTS:

Morantel Tartrate: 0.194% (0.88 grams per pound)

INGREDIENTS:

Wheat middlings, soybean hulls (max 20%), calcium carbonate, molasses products, calcium propionate (a preservative).

DIRECTIONS FOR USE:

- Feed as a single therapeutic treatment at 0.44 grams of morantel tartrate per 100 pounds of body weight.
- Fresh water should be available at all times.
- After medicated feed is consumed, resume normal feeding.
- Conditions of constant worm exposure may require retreatment in 2 to 4 weeks.
- When used as a top dress the medication as well as the underlying feed should be evenly distributed.

WARNING:

Do not treat goats within 30 days of slaughter. No milk discard required following use in dairy goats. For use in goats only.

ISSU15XB11
Distributed by Durvet, Inc.
Blue Springs, MO 64014
www.durvet.com | 1-800-821-5570

NET WT: 3 LB (1.36 KG)

GUARANTEED ANALYSIS:

Crude Protein (min)	12%
Crude Fat (min)	1%
Crude Fiber (max)	15%
Acid Detergent Fiber (max)	23%
Calcium (min)	2.8%
Calcium (max)	3.3%
Phosphorus (min)	0.5%
Selenium (min)	0.2 PPM

TOP-DRESS DIRECTIONS:

Body Weight (LB)	GoatCare•2X
12.5	1 oz
25	2 oz
50	4 oz
75	6 oz
100	8 oz
150	12 oz
200	16 oz

4 oz MEASURING SCOOP ENCLOSED.
1 Level Scoop per 50 lb. Body Weight

CAUTION:

Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment and control of parasitism. Do not mix in feeds containing Bentonite.

