



Pyrantel•S

TYPE C MEDICATED FEED

SWINE ANTHELMINTIC

Contains Banminth®(Pyrantel Tartrate)

TREATS
5
FORTY LB
PIGS

For removal and control of
large roundworm (*Ascaris suum*) and
nodular worm (*Oesophagostomum*)
infections in swine.

ACTIVE DRUG INGREDIENT

Pyrantel Tartrate 0.0881% (800 grams per ton)

Banminth is a registered trademark of Phibro Animal Health Corporation for Pyrantel Tartrate.



See Back Panel for Additional Information

NET WT: 5 LB (2.26 KG)



Pyrantel•S

TYPE C MEDICATED FEED

SWINE ANTHELMINTIC

For removal and control of
 large roundworm (*Ascaris suum*) and
 nodular worm (*Oesophagostomum*) infections in swine.



ACTIVE DRUG INGREDIENT:

Pyrantel Tartrate: 0.0881% (800 grams per ton)

INGREDIENTS:

Grain products, processed grain by-products, plant protein products, calcium carbonate, sodium chloride, monocalcium phosphate, L-Lysine, roughage products, iron sulfate, zinc sulfate, copper sulfate, manganous oxide, dried *Pichia pastoris* fermentation extract, vitamin E supplement, niacin supplement, d-calcium pantothenate, sodium selenite, vitamin A acetate, riboflavin supplement, menadione nicotinamide bisulfite (source of vitamin K and niacin), vitamin B12 supplement, vitamin D₃ supplement and ethylenediamine dihydriodide.

CAUTION:

Consult your veterinarian before using in severely debilitated animals. Do not mix in feeds 24 hours prior to slaughter.

WARNING:

Withdraw this feed 24 hours prior to slaughter.

The SAFETY DATA SHEET (SDS) contains detailed occupational safety information and is available upon request at 1-800-821-5570.

GUARANTEED ANALYSIS:

Crude Protein (min)	16%
Lysine (min)	0.90%
Crude Fat (min)	3.00%
Crude Fiber (max)	6.00%
Calcium (min)	0.60%
Calcium (max)	0.70%
Phosphorus (min)	0.60%
Salt (min)	0.30%
Salt (max)	0.40%
Selenium (min)	0.30 PPM
Zinc (min)	165 PPM

DIRECTIONS FOR USE:

- Enclosed scoop measures approximately 0.50 pound (8 oz).
- Feed as the sole ration for a single therapeutic treatment at the rate of 1 pound of feed per 40 pounds of body weight for animals up to 200 pounds. For animals 200 pounds or over, feed 5 pounds of feed per head. Pigs will generally consume this feed within a few hours, after which regular feeding should be resumed.
- Better worming is achieved when pigs are separated by size into different lots or pens for treatment.
- Water should be made available to animals during fasting and treatment periods.
- Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

ISSU16XB03

Distributed by Durvet, Inc.

Blue Springs, MO 64014

www.durvet.com | 1-800-821-5570

NET WT: 5 LB (2.26 KG)

